


EXPLANATORY DATA  
LOS ANGELES SHEET  
GEOLOGIC MAP OF CALIFORNIA

OLAF P. JENKINS EDITION

Compiled by Charles W. Jennings and Rudolph G. Strand, 1969

THIS DATA SHEET IS A REPRINT OF THE DATA SHEET ACCOMPANYING THE LOS ANGELES SHEET, GEOLOGIC MAP OF CALIFORNIA, OLAF P. JENKINS EDITION, FIRST PUBLISHED IN 1969. IT HAS NOT BEEN ALTERED. THE GEOLOGY SHOWN ON THE LOS ANGELES SHEET OF THE BOUGUER GRAVITY MAP OF CALIFORNIA IS ALSO REPRINTED FROM THE GEOLOGIC MAP OF CALIFORNIA, 1969. THE GRAVITY DATA PRESENTED WERE COMPILED IN 1974 AND PUBLISHED IN 1975.

INDEX TO GEOLOGIC MAPPING  
USED IN THE COMPILATION OF THE LOS ANGELES SHEET



1. Baddley, E. R., 1954, Geology of the South Mountain oil field, Ventura County, California in *Geology of Southern California*; California Div. Mines Bull. 170, Map Sheet 29, scale 1 inch = approx. 1550 feet.
2. Bailey, Thomas L. (consulting geologist, Ventura, Calif.), Geology of a portion of Ventura basin, Los Angeles & Ventura Counties, California, scale 1:24,000, unpublished, 1951. (Modification and additions to the area west of 119°00' from unpublished mapping and compilation by T. W. Dibblee, Jr. Modifications and additions in the Hopper Mountain and lower Piru Canyon areas by W. H. Corey and by Paul Siemon, Continental Oil Company 1946. East of 119° principal published reference: Kew, William S. W., 1924, Geology and oil resources of a part of Los Angeles and Ventura Counties, California: U. S. Geol. Survey Bull. 753, 202 p., Pl. I, scale 1:62,500.)
3. Bain, Roland J., Geology of the Eureka Canyon area, Ventura County, California, scale 1:12,000, Univ. California, Los Angeles, unpublished M.A. thesis, 1954.
4. Bishop, William C., Geology of the southern flank of Santa Susana Mountains, county line to Limekiln Canyon, Los Angeles County, California, scale 1:12,000, Univ. California, Los Angeles, unpublished M.A. thesis, 1950.
5. Black, Bruce A., Geologic map of Topatopa area, scale 1:250,000, unpublished mapping, Shell Oil Company.
6. Blackerby, Bruce A., The Conejo volcanics in the Malibu Lake area of the western Santa Monica Mountains, Los Angeles County, California, scale 1:12,000, Univ. California, Los Angeles, unpublished PhD thesis, 1955.
7. Brady, Thomas J., Geology of part of the central Santa Monica Mountains, east of Topanga Canyon, Los Angeles County, California, scale 1:12,000, Univ. California, Los Angeles, unpublished M.A. thesis, 1957.
8. Brown, George E., Geology of parts of the Calabasas and Thousand Oaks quadrangles, Los Angeles and Ventura Counties, California, scale 1:18,000, Univ. California, Los Angeles, unpublished M.A. thesis, 1957.
9. Burnett, John L., The geology of the southern portion of Frazier Mountain near Gorman, California, scale 1:31,680 and 1:7,920, Univ. California, Berkeley, unpublished M.S. thesis, 1960. Crowell, John C., 1964 (see ref. 18).
10. Byer, John W., Geologic map of a part of the San Rafael Hills, scale 1:24,000, unpublished map for City of Glendale, 1968.
11. California Dept. Water Resources, 1966, Planned utilization of ground water basins—San Gabriel Valley: Bull. 104-2. Appendix A: Geohydrology, Pl. 9A, scale 1:125,000. Buwalda, John P., Geologic map of Raymond basin prepared for Pasadena Water Dept., 1940, scale 1:24,000, unpublished.
12. Carman, Max F., Jr., 1964, Geology of the Lockwood Valley area, Kern and Ventura Counties, California: California Div. Mines and Geology Special Rept. 81, 62 pp., Pl. 1, scale approx. 1:31,680.
13. Carter, Neville L., Geology of the Fernwood-Topanga Park area, Santa Monica Mountains, California, scale 1:12,000, Univ. California, Los Angeles, unpublished M.A. thesis, 1958. Hazenbush, George C., Geology of the eastern part of the Dry Canyon and Las Flores quadrangles, Los Angeles County, California, scale 1:12,000, Univ. California, Los Angeles, unpublished M.A. thesis, 1950.
14. Castle, Robert O., 1960, Surficial geology of the Beverly Hills and Venice quadrangles, California: U. S. Geological Survey open file report, map scale 1:24,000. Castle, Robert O., 1960, Geology of the Baldwin Hills area, California: U. S. Geol. Survey open file report, map scale 1:12,000. (Northwest trending concealed faults based on Poland, et al, (ref. 66). Santa Monica fault from John E. Kilkenny, Union Oil Co. of California, written communication 7/12/68.)
15. Champeny, Jon D., Paleocene and Upper Cretaceous stratigraphy of Santa Ynez Canyon and adjacent areas, Santa Monica Mountains, California, scale 1:24,000, Univ. California, Los Angeles, unpublished M.A. thesis, 1961.
16. Conrad, Stanley D., Geology of the eastern portion of the Simi Hills, Los Angeles and Ventura Counties, California, scale approx. 1:30,000, Univ. California, Los Angeles, unpublished M.A. thesis, 1949.
17. Corbató, Charles E., 1963, Bouguer gravity anomalies of the San Fernando Valley, California: Univ. California Dept. Geol. Sci. Bull., v. 46, no. 1, pp. 1-32, Map 1, scale 1:48,000 and fig. 7, scale 1 inch = approx. 5 miles. State (California) Water Rights Board, 1962, San Fernando Valley reference report of referee (City of Los Angeles vs. City of San Fernando, et al., No. 650079 Superior Court, Los Angeles County) v. 1, 258 pp., Pl. 4, scale 1 inch = approx. 2.5 miles. Eckis, Rollin, 1934, Geology and ground water storage capacity of valley fill: California Div. Water Resources Bull. 45, 279 pp., Pl. A, scale 1 inch = approx. 2.25 miles.
18. Crowell, John C., 1964, The San Andreas fault zone from the Temblor Mountains to Antelope Valley, southern California: in Pacific Section A.A.P.G.-S.E.P.M. and San Joaquin Geol. Soc. Guidebook, pp. 8-38, "Pl. 1", by John C. Crowell and others, scale 1:62,500 (Plush Ranch—Caliente Fm. contact northwest of San Guillermo Mtn. from Kahle, James E., Megabreccias and sedimentary structures of the Plush Ranch Formation, northern Ventura County, California, Pl. 3, scale 1:62,500, Univ. California, Los Angeles, unpublished M.A. thesis, 1966).
19. Crowell, John C., et al., Geologic map of Ridge basin, southern California, scale 1:24,000, faculty research Univ. California, Los Angeles, unpublished 1956 (with later revision by J.C.C.). Crowell, John C., 1954, Geology of the Ridge basin area, Los Angeles and Ventura Counties: California Div. Mines Bull. 170, Map sheet 7, scale 1 inch = approx. 1.5 miles. Crowell, John C., 1952, Geology of the Lebec quadrangle, California: California Div. Mines Spec. Rept. 24, 23 p., Pl. 1, scale 1:31,680.
20. Crowell, John C., Geology of the Santa Felicia—Canton Canyon area, Los Angeles and Ventura Counties, California, scale 1:14,400, faculty research, University California, Los Angeles unpublished 1953. (Area covered in part by Kříž, Stanislav J., Stratigraphy and structure of the Whitaker Peak—Reasoner Canyon area, Ventura and Los Angeles Counties, California, scale 1:24,000, Princeton Univ., unpublished Ph.D. thesis, 1947.)
21. Crowell, John C., McClure, D. V., Kienle, C. F., and Edwards, L. N., Geologic map of the eastern end of the Santa Ynez fault zone, scale 1:250,000, Univ. California, Santa Barbara, unpublished field studies, 1967.
22. Daviss, Steven N. and Woodford, A. O., 1949, Geology of the northwestern Puente Hills, Los Angeles County, California: U. S. Geol. Survey Oil and Gas Prelim. Map 83, Sheet 1 of 2, scale 1:12,000.
23. Dibblee, Thomas W., Jr., 1960, Geologic map of the Lancaster quadrangle, Los Angeles County, California: U. S. Geol. Survey Map MF-76, scale 1:62,500. Wallace, Robert E., 1949, Structure of a portion of the San Andreas rift in southern California: Geol. Soc. America Bull. v. 60, p. 781-806, Pl. 1, scale 1:31,680. Noble, Levi F., 1953, Geology of the Pearland quadrangle, California: U. S. Geol. Survey GQ 24, scale 1:24,000.
24. Dibblee, Thomas W., Jr., 1961, Geologic map of the Bouquet Reservoir quadrangle, Los Angeles County, California: U. S. Geol. Survey Map MF-79, scale 1:62,500. Wallace, Robert E., 1949, Structure of a portion of the San Andreas rift in southern California: Geol. Soc. America Bull. v. 60, p. 781-806, Pl. 1, scale 1:31,680.
- 24b. Dibblee, Thomas W., Jr., 1961, (see ref. 24a). Muehlberger, William R., 1958, Geology of northern Soledad basin, Los Angeles County, California: Am. Assoc. Petroleum Geologists Bull., v. 42, no. 8, pp. 1812-1894, Fig. 3, scale 1 inch = approx. 4,300 feet.
25. Dibblee, Thomas W., Jr., 1961, Map in Geology and paleontology of the southern border of the San Joaquin Valley, Kern County, California: Pacific Section S.E.P.M.-S.E.G. A.A.P.G. and San Joaquin Geol. Soc. Guidebook, Pl. II, scale approx. 1:62,500.
26. Dibblee, Thomas W., Jr., 1963, Geology of the Willow Springs and Rosamond quadrangles, California: U. S. Geol. Survey Bull. 1089-C, 253 pp., Pl. 10, scale 1:62,500.
27. Dibblee, Thomas W., Jr., Geology of the Tecuya Creek and Pastoria Creek quadrangles, scale 1:31,680, unpublished, 1955. Dibblee, Thomas W., Jr., Bruer, W. G., Hackel, Otto, Warne, A. H., 1965, Map in Geology of southeastern San Joaquin Valley, California: Pacific Section A.A.P.G.-S.E.G.-S.E.P.M. Guidebook, Map, scale 1:48,000.
28. Dibblee, Thomas W., Jr., 1966, Geology of the central Santa Ynez Mountains, Santa Barbara County, California: California Div. Mines and Geology Bull. 186, 99 pp., Pl. 1, scale 1:31,680, and Pl. 3, scale 1:62,500 (additions from J. G. Vedder and others ref. 87).
29. Dibblee, Thomas W., Jr., Geology of part of the Morro Hill and Reyes Peak quadrangles, California, scale 1:31,680, unpublished, 1946 (modified in part by information from Stanford Geological Survey ref. 78).
30. Dibblee, Thomas W., Jr., Geologic map of the McPherson Peak quadrangle, California, scale 1:62,500, unpublished, 1948-49, 1962.
31. Dibblee, Thomas W., Jr., Geology of the San Guillermo, Lockwood Valley, Lion Canyon, and Topatopa Mountains quadrangles, California, scale 1:62,500, unpublished, 1949 (modified in part by Crowell, and Kahle, see ref. 18).
32. Dibblee, Thomas W., Jr., Geologic maps of parts of the Quail, Manzanita, and Liebre quadrangles, California, scale 1:24,000, unpublished, 1957. Dibblee, Thomas W., Jr., 1967, Areal geology of the western Mojave Desert, California: U. S. Geol. Survey, Prof. Paper 522, Pl. 1, scale 1:125,000, and figs. 35 and 36, scale approx. 1:62,500.
33. Dibblee, Thomas W., Jr., Geologic map of a part of the Salisbury Canyon quadrangle, California, scale 1:62,500, unpublished, 1962.
34. Dibblee, Thomas W., Jr., Geologic map of part of the Acton quadrangle, California, scale 1:62,500, unpublished, 1968.
35. Dibblee, Thomas W., Jr. and Fisher, Richard V., Geology of the Wheeler Springs and Old Man Mountain quadrangles, California, scale 1:31,680, unpublished, 1946 and 1958. Fisher, Richard V. and Dibblee, Thomas W., Jr., 1961, Geology and possible tectonic significance of Munson Creek fault, San Rafael Mountains, California: Amer. Assoc. Petroleum Geol. Bull. v. 45, no. 9, pp. 1572-1581.
36. Dibblee, Thomas W., Jr. (partly compiled from other sources). Geologic map of the "Ventucopa" quadrangle, California, scale 1:62,500, unpublished, 1946-66.
37. Dibblee, Thomas W., Jr. (partly compiled from other sources). Geologic map of the Ventura quadrangle, California, scale 1:62,500, unpublished 1930-45. Standard Oil Company of California, Western Operations, Inc., Geologic map D2873T, scale 1:24,000, unpublished.
38. Ehlig, Perry L., Geologic map of a part of the central San Gabriel Mountains, California, scale 1:250,000, California State College at Los Angeles, unpublished faculty research, 1967. (Additions by T. W. Dibblee, Jr., Geologic maps of part of the Condor Peak, Chilao Flat, Pasadena and Mt. Wilson quadrangles, scale 1:24,000, U. S. G. S. work in progress, 1967-68.)
39. Elam, Jack G., Geology of the Seminole quadrangle, Los Angeles County, California, scale 1:20,000, Univ. California, Los Angeles, unpublished M.A. thesis, 1948.

(Continued on back page)

# STRATIGRAPHIC NOMENCLATURE—LOS ANGELES SHEET

AGE	STATE MAP SYMBOL	STATE MAP UNIT <small>State Map Units listed here are not necessarily in stratigraphic sequence; the sequence used has been standardized for all sheets of the Geologic Map of California</small>	STRATIGRAPHIC UNITS AND CHARACTERISTIC LITHOLOGIES <small>Formally named formations grouped in sequence (separated by semicolons) are listed from youngest to oldest.</small>			
QUATERNARY	Recent	Qs	<b>RECENT DUNE SAND</b>	Wind-blown sand and beach deposits along coast, desert dune sand and bars of wave-deposited sand around Rosamond Dry Lake.		
		Ql	<b>QUATERNARY LAKE DEPOSITS</b>	Playa clay and silt of Rosamond Dry Lake.		
		Qal	<b>RECENT ALLUVIUM</b>	"Younger" alluvium consisting of Recent clay, silt, sand and gravel, unconsolidated, poorly stratified to well stratified, includes alluvial fan, flood-plain, and streambed deposits. In some desert areas includes mixture of playa clay and wind-blown sand.		
		Qf	<b>RECENT ALLUVIAL FAN DEPOSITS IN THE GREAT VALLEY</b>	Alluvial fan deposits at the southern end of the San Joaquin Valley.		
		Qt	<b>QUATERNARY NONMARINE TERRACE DEPOSITS</b>	Alluvial terrace deposits, in places along the coast overlies thin marine terrace deposits. Locally includes fanglomerate and "older" alluvial debris at higher levels.		
	Pleistocene	Pliocene	Qc	<b>PLEISTOCENE NONMARINE SEDIMENTARY DEPOSITS</b>	Casitas Formation—red to greenish clay, silt, sand and conglomerate (Carpinteria area). Frazier Mountain Formation—conglomerate, fanglomerate and landslide breccia of local derivation (Lockwood Valley). Pacoima Formation—dark-brown breccia and fanglomerate, locally folded. Nadeau Gravel—gravel and interbedded sand of prevailing dark color (Pearland quadrangle). Harold Formation—interbedded gravel, sand, silt, and gypsiferous clay. "Older" alluvium consisting of gravel, sand, clay, and silt; dissected alluvial fan deposits. "Ancient" alluvium, commonly with soil formation on surface (San Gabriel Valley).	
			Qm	<b>PLEISTOCENE MARINE DEPOSITS AND MARINE TERRACE DEPOSITS</b>	San Pedro Formation—sand, silt, and marl, highly fossiliferous (Los Angeles basin). Marine and some stream terrace deposits along the coast and on Santa Cruz Island. Qm-Pu: Santa Barbara Formation—buff to yellow sand, silt and calcareous sediments.	
			QP	<b>PLIOCENE-PLEISTOCENE NONMARINE SEDIMENTARY DEPOSITS</b>	Paso Robles Formation—interbedded gray to brown clay, sandstone, and conglomerate (Cuyama area and Santa Ynez Mountains). Saugus Formation—sandstone, conglomerate and siltstone, mainly fluvial but locally brackish water deposits. Tulare Formation—gravel, clay and conglomerate (south end San Joaquin Valley).	
			Puc	<b>UNDIVIDED UPPER PLIOCENE NONMARINE SEDIMENTARY ROCKS</b>	Hungry Valley Formation—brown conglomerate, white conglomeratic sandstone, and brown mudstone.	
			Pc	<b>UNDIVIDED PLIOCENE NONMARINE SEDIMENTARY ROCKS</b>	Morales Formation—gray, poorly bedded stream-laid gravel and sand; Quatal Formation—orange to buff, pebbly, arkosic sandstone and light-red clays, gypsiferous siltstone, and gypsum; Lockwood Clay of Carman, 1964—pale-tan to gray montmorillonitic clay and interbedded fine-grained sandstone. Peace Valley Formation—brown siltstone, buff sandstone, some conglomerate and sedimentary breccia. Meeke Mine Formation—pebble gravel, sand, and lacustrine clay shale; may be early Pleistocene in age (Tehachapi Mountains area).	
			Pu	<b>UPPER PLIOCENE MARINE SEDIMENTARY ROCKS</b>	Sunshine Ranch Member of Saugus Formation—continental and brackish-water greenish sandstone and mudstone, red conglomerate beds, and thin limestone beds (San Fernando Reservoir area); upper member of the Fernando Formation (includes rocks commonly called Pico Formation)—sandy siltstone, conglomerate and sandstone (Los Angeles basin). Pico Formation—siltstone, sandstone and conglomerate (Ventura basin). Careaga Sandstone—poorly consolidated, fine- to medium-grained sandstone (Santa Ynez Mtns.). Unnamed marine Pliocene-Pleistocene (?) silt and sand in the Baldwin Hills area.	
			Pmlc	<b>MIDDLE AND/OR LOWER PLIOCENE NONMARINE SEDIMENTARY ROCKS</b>	Anaverde Formation—gray shale, arkosic sandstone, red conglomerate, dark-gray dioritic breccia; plant fossils of early to middle Pliocene age (in San Andreas fault zone). Ridge Route Formation—sandstone with conglomerate, shale, and siltstone interbeds. Chanac Formation—sandstone, claystone and conglomerate.	
			Pml	<b>MIDDLE AND/OR LOWER PLIOCENE MARINE SEDIMENTARY ROCKS</b>	Lower member of the Fernando Formation (includes rocks commonly called Repetto Formation)—siltstone, conglomerate and fine sandstone. Etchegoin Formation—buff, fine, fossiliferous sand (SW San Joaquin Valley). Unnamed siliceous mudstone in the Bitterwater Creek area (may be late Miocene in age). Towsley Formation—siltstone, mudstone, sandstone and conglomerate (late Miocene and early Pliocene age; Santa Susanna Mountains).	
			Mc	<b>UNDIVIDED MIOCENE NONMARINE SEDIMENTARY ROCKS</b>	Fiss Fanglomerate—fanglomerates of brown volcanic detritus and gray granitic detritus (Miocene age uncertain; Willow Springs and Rosamond quadrangles). Punchbowl Formation—white, buff to pink sandstone, gray to red siltstone and clay shale, and gray to red conglomerate (middle and late Miocene and early Pliocene age; Valyermo area). Mint Canyon Formation—grayish-green or light-brown siltstone and mudstone, light-brown, gray or reddish sandstone and conglomerate; some tuff beds (late Miocene and early Pliocene age; Mint Canyon area). Oso Canyon Formation—fanglomerate, gray-white to red sandstone, and red to green siltstone, grades laterally west into Quail Lake Formation (late Miocene age; west end Antelope Valley). Bissell Formation—dolomite, limestone and shale with magnesite layers, sandstones and conglomerates (Miocene or Pliocene age; Bissell Hills, Rosamond quadrangle). Unnamed gray fanglomerate sand and clay that grades northward into marine sandstone and shale of middle and late Miocene age (southern San Joaquin Valley).	
			Mu	<b>UPPER MIOCENE MARINE SEDIMENTARY ROCKS</b>	Tequepis Sandstone—gray-white, massive to thick-bedded, compact, semifriable sandstone (Santa Ynez Mtns.). Sisquoc Formation—soft, fissile to massive diatomite in part tuffaceous, shale and siltstone (Santa Ynez Mtns.). Santa Margarita Formation—sandstone, siltstone, some conglomerate (Cuyama Valley and Pine Mtn. areas). Modelo Formation—siliceous and diatomaceous shale, sandstone, siltstone and some conglomerate (in part middle Miocene; Santa Monica Mtns.). Upper part of Monterey Shale on Pt. Dume. Castaic Formation—shale with interbedded sandstone and beds of pebble conglomerate (considered to be same as the Modelo Formation by some geologists). Puente Formation—siltstone, siliceous shale, diatomaceous shale, sandstone and conglomerate, including Sycamore Canyon Member—conglomerate and siltstone (Los Angeles area). Quail Lake Formation—buff sandstone and brown shale (west end Antelope Valley).	
Mmc	<b>MIDDLE MIOCENE NONMARINE SEDIMENTARY ROCKS</b>	Tick Canyon Formation—siltstone, sandstone, conglomerate, probably early Miocene. Unnamed nonmarine sandstone, mudstone and conglomerate (lateral equivalent to part of lower Topanga Formation of Durrell, 1954, Santa Monica Mtns.). Unnamed cobble conglomerate and red beds (southern San Joaquin Valley).				
Mm	<b>MIDDLE MIOCENE MARINE SEDIMENTARY ROCKS</b>	Monterey Shale—brown, soft, fissile, punky, organic shale, hard porcellaneous shale and chert, bituminous silty siliceous shale, locally includes thin limestone beds, siliceous clay siltstone and sandstone; south of Malibu Coast fault includes shale, sandstone, dolomite, and chert. Unnamed sandstone, mudstone, sedimentary breccia and some volcanic rocks (south of Malibu Coast fault). San Onofre Breccia—sandstone and breccia of schist fragments including slabs and blocks of glaucophane schist (along coast, western Santa Monica Mtns.). Topanga Formation—shale, sandstone, some conglomerate, arkose (Santa Monica Mtns.).				
TERTIARY	Miocene	Mv	<b>MIOCENE VOLCANIC ROCKS: UNDIFFERENTIATED</b>	Interbedded agglomerate, flow breccias, flows, tuffs, and volcanic sandstone (central Santa Monica Mtns.). Andesite and basalt flows in the Topanga Formation(?), Verdugo Mtns. and Pacoima Hills.		
		Mv <sup>a</sup>	<b>ANDESITIC</b>	Conejo Volcanics—hypersthene andesite (Santa Monica Mtns.).		
		Mv <sup>b</sup>	<b>BASALTIC</b>	Conejo Volcanics—basalt flows and breccia, some andesite, arkose and tuff (Santa Monica Mtns.). Vesicular and porphyritic extrusive basaltic flows and pyroclastic rocks (Conejo Volcanics?; Santa Cruz and Anacapa Islands). Basalt flows in the Gem Hill Formation (Miocene age uncertain; Rosamond quadrangle). Vesicular amygdaloidal basalt flows, with interbedded tuff breccia and agglomerate (Sunland area).		
		Mv <sup>p</sup>	<b>PYROCLASTIC</b>	Gem Hill Formation—mainly tuff, tuff breccia, and tuffaceous sandstone (Miocene age uncertain) and Bobtail Quartz Latite Member—felsite breccia (Rosamond-Willow Springs area). Conejo Volcanics—agglomerate, pyroclastic and interbedded epiclastic rocks (Santa Monica Mtns.). Rhyolite, andesite, and basalt breccia, agglomerate, scoriaceous flows, tuff, and tuff breccia (Santa Cruz Island).		
		MI	<b>LOWER MIOCENE MARINE SEDIMENTARY ROCKS</b>	"Temblor" Sandstone—arkosic sandstone, pebbly in part, calcareous in part (Santa Ynez River area); Rincon Shale—blue-gray, silty, micaceous shale (Santa Barbara-Ojai area); Vaqueros Formation—sandstone, siltstone, some conglomerate; in the Caliente Range divided into: Painted Rock Sandstone Member; Soda Lake Shale Member; Soda Lake Sandstone Member. Pleito Formation—sandstone and interbedded clay shale (in part Oligocene). Unnamed sandstone and conglomerate on Santa Cruz Island, including glaucophane schist-bearing breccias and conglomerates "San Onofre Breccia" of Rand, 1931, ref. no. 80 (similar breccias occur on Anacapa Island where shown as Mv <sup>b</sup> ). Unnamed brown to white sandstone, gray claystone and siltstone, in part middle Miocene (San Rafael Mtns. area).		
		Oligocene	Φc	<b>OLIGOCENE NONMARINE SEDIMENTARY ROCKS</b>	Sespe Formation—red to gray-green shale, sandstone, conglomerate, and breccias (probably ranges in age from late Eocene to early Miocene). Vasquez Formation—red to light-brown arkosic sandstone and conglomerate, some interbedded maroon siltstone and breccias (Oligocene or early Miocene age). Simmler Formation (Plush Ranch Formation of Carman)—red, green, and bluish-gray clay, arkosic sandstone, conglomerate, fanglomerate, and breccia (Oligocene or early Miocene age; Cuyama-Lockwood Valley areas). Tecuya Formation—red and green conglomerate sandstone and clay (early Miocene and Oligocene(?) age; south end San Joaquin Valley). Unnamed, greenish-gray to red conglomerate sandstone and mudstone of Miocene or Oligocene age (San Rafael Mountains).	
			Φ	<b>OLIGOCENE MARINE SEDIMENTARY ROCKS</b>	Caviota Formation—massive to thick-bedded, buff-weathering arkosic sandstone. "San Emigdio" Formation—alternate beds of yellowish and grayish sandstones and shales (San Emigdio Mtns.).	
			Oligocene Volcanic Rocks:	Φv	<b>UNDIFFERENTIATED</b>	Volcanic rocks of the Vasquez Formation, undifferentiated—black and reddish vesicular and amygdaloidal basalt and andesite, purplish breccia, minor beds of lithic tuff; associated with lacustrine strata in Tick Canyon containing borate beds. Volcanic rocks in the Tecuya Formation—olivine basalt, and andesite flows.
				Φv <sup>r</sup>	<b>RHYOLITIC</b>	Neenach Volcanic Formation—light-gray to pink rhyolitic felsite, Oligocene(?) and early or possibly middle Miocene.
				Φv <sup>a</sup>	<b>ANDESITIC</b>	Volcanic rocks of the Vasquez Formation, andesitic—dark reddish-brown andesite flow breccias, including some greenish-white to buff tuff breccias. Neenach Volcanic Formation—dark-brown andesite, Oligocene(?) and early or possibly middle Miocene.
Φv <sup>b</sup>	<b>BASALTIC</b>	Volcanic rocks of the Vasquez Formation, basaltic—greenish-black basalts, commonly amygdaloidal. Volcanic rocks of the Plush Ranch Formation—black or red-brown hypersthene-augite-olivine basalt; some tuff and associated lacustrine strata that contain borate deposits. (Carman, ref. 12.)				
E	<b>EOCENE MARINE SEDIMENTARY ROCKS</b>	"Coldwater" Sandstone—arkosic sandstone, local oyster reefs, siltstone and shale; Sacate Formation—argillaceous to silty shale, some sandstone; Cozy Dell Shale—argillaceous to silty shale; Matilija Sandstone—arkosic sandstone; Anita Shale—shale and some sandstone; Juncal Formation—shale and arkosic sandstone; Sierra Blanca Limestone—massive, hard, organic limestone (Santa Ynez, Topatopa and San Rafael Mtns.); Tejon Formation—sandstone, siltstone, shale, some conglomerate (San Emigdio-Tehachapi Mtns.). Lajas Formation—conglomerate, sandstone, siltstone; Santa Susana Formation—concretionary shale, sandstone, conglomerate lenses (Paleocene in part). Unnamed rocks of middle or late Eocene age in Elsmere Canyon. Unnamed Eocene siltstones, sandstones, shales and conglomerates in northern San Rafael Mountains, Santa Cruz Island, Piru Creek and Elizabeth Lake Canyon areas.				

# STRATIGRAPHIC NOMENCLATURE — Continued

AGE	STATE MAP SYMBOL	STATE MAP UNIT <small>State Map Units listed here are not necessarily in stratigraphic sequence; the sequence used has been standardized for all sheets of the Geologic Map of California</small>	STRATIGRAPHIC UNITS AND CHARACTERISTIC LITHOLOGIES <small>Formally named formations grouped in sequence (separated by semicolons) are listed from youngest to oldest.</small>	
CENOZOIC	TERTIARY	Undivided	Paleocene	
			Paleocene	
		Ep	<b>PALEOCENE MARINE SEDIMENTARY ROCKS</b>	Pattway Formation— <i>siltstone, locally concretionary, sandstone, pebble-boulder conglomerate</i> . "Martinez" Formation— <i>sandstone, conglomerate, mudstone</i> (includes some rocks of Eocene age); Las Virgenes Sandstone— <i>gray sandstone</i> (Simi Hills); Simi Conglomerate— <i>well rounded pebbles and cobbles up to 1 foot diameter in matrix of coarse-grained arkosic sandstone; lenticular beds of sandstone</i> . Unnamed sandstone interbedded with thin shale and conglomerate (Paleocene; Santa Cruz Island, Santa Monica Mtns.). Unnamed conglomerates, sandstones, and siltstones (Paleocene and lower Eocene?; Castaic Creek-Elizabeth Canyon area). Ep-E: San Francisquito Formation— <i>arkosic sandstone, conglomerate and shale</i> . (Paleocene and possibly Eocene, stratigraphically above the unnamed conglomerates, etcetera, described above.)
		Tc	<b>TERTIARY NONMARINE SEDIMENTARY ROCKS</b>	Caliente Formation— <i>calcareous, arkosic sandstone, red and gray mudstone, channel sand, and conglomerate; reddish-brown, thick-bedded, argillaceous, coarse-grained sandstone</i> (now dated as early Miocene to middle Pliocene). Violin Breccia— <i>rubble of gneiss blocks up to 6 feet in diameter in a mudstone matrix, accumulated as talus or alluvial debris at the base of the San Gabriel fault scarp; grades into finer grained Ridge Route Formation to the east</i> . (Miocene to Pliocene or Pleistocene age.)
		Tm	<b>TERTIARY MARINE SEDIMENTARY ROCKS</b>	Unnamed shale and sandstone, some conglomerate, travertine, and tuff (except for travertine and arkose, has a marine aspect; age uncertain, may be Eocene, T. W. Dibblee, Jr., written communication, 1968; located at intersection of Big Pine and San Andreas faults).
		Tv <sup>b</sup>	<b>TERTIARY VOLCANIC ROCKS:</b> <b>BASALTIC</b>	Basalt flows of the Caliente Formation— <i>multiple olivine basalt flows, highly vesicular in part, includes breccias on some flow tops</i> (Miocene age).
		Ti	<b>TERTIARY INTRUSIVE (HYPABYSSAL) ROCKS:</b> <b>UNDIFFERENTIATED</b>	Intrusive rocks ranging from olivine basalt to dacite; middle Miocene(?) age (Santa Monica Mtns.).
		Ti <sup>r</sup>	<b>RHYOLITIC</b>	Bobtail Quartz Latite Member of Gem Hill Formation— <i>intrusive facies including felsite and porphyritic facies, porphyry and perlite</i> ; Miocene(?) age; Rosamond-Willow Springs area. Rhyolite and obsidian plugs (Frazier Mountain area).
		Ti <sup>a</sup>	<b>ANDESITIC</b>	Andesite porphyry of Miocene(?) or Oligocene(?) age (San Emigdio Mtns.). Hornblende andesite sill (South Mountain).
		Ti <sup>b</sup>	<b>BASALTIC</b>	Intrusive basalt and diabase of middle Miocene(?) age (Santa Monica Mtns.).
CRETACEOUS		Ku	<b>UPPER CRETACEOUS MARINE SEDIMENTARY ROCKS</b>	Ku: Jalama Formation— <i>gray to black shale, hard, light-gray arkosic sandstone, and cobble conglomerate</i> (Santa Ynez Mtns.). "Chico" Formation— <i>sandstone, shale and conglomerate</i> . Unnamed light-gray sandstone, dark-gray siltstone and shale, and minor cobble conglomerate. Ku: Local lenses of presumably non-marine greenish- to reddish-gray conglomerate, sandstone, and mudstone (San Rafael and Santa Susana Mtns.).
		Kl	<b>LOWER CRETACEOUS MARINE SEDIMENTARY ROCKS</b>	Espada Formation— <i>dark olive-gray shale, siltstone, and thin sandstone beds</i> (San Rafael and Santa Ynez Mtns.).
		KJf	<b>FRANCISCAN FORMATION</b>	Franciscan Formation— <i>sheared and severely deformed, dark greenish-gray graywacke, black claystone and chert</i> (San Rafael and Santa Ynez Mtns.).
		KJfv	<b>FRANCISCAN VOLCANIC AND METAVOLCANIC ROCKS</b>	Greenstone of the Franciscan Formation and minor quantities of glaucophane schist (San Rafael and Santa Ynez Mtns.).
		gr	<b>MESOZOIC GRANITIC ROCKS</b> <b>UNDIFFERENTIATED</b>	Granite and other quartz-bearing plutonic rocks (Soledad basin). Biotite granodiorite, quartz monzonite, and quartz diorite (Griffith Park). Areas of undivided, predominately granitic rocks; locally may grade into rocks of banded "gneissic" structure (gneissic granite) or may be complexly associated with gneiss (Sawmill Mountain and Bouquet Reservoir areas). Mount Lowe "Granodiorite" and Parker "Quartz Diorite"— <i>light-colored, foliated, quartz-poor, feldspar-rich, porphyritic granitic rock of monzonitic to dioritic composition</i> (Permian-Triassic; San Gabriel Mtns.).
MESOZOIC		gr <sup>a</sup>	<b>ADAMELLITE (QUARTZ MONZONITE), GRANITE, ALASKITE</b>	Liebre Quartz Monzonite— <i>fractured, gray hornblende-biotite quartz monzonite, with dark inclusions and pegmatites</i> . Lebec Quartz Monzonite— <i>gray hornblende-biotite quartz monzonite</i> . Mt. Pinos Granite— <i>biotite granite and hornblende-biotite quartz diorite</i> . Tejon Lookout Granite— <i>biotite granite</i> (Tehachapi Mtns.). Unnamed granites and quartz monzonites (north of San Andreas fault).
		gr <sup>g</sup>	<b>GRANODIORITE</b>	Unnamed granodiorite including some quartz diorite, quartz monzonite, and pink granite (San Gabriel Mtns.).
		gr <sup>t</sup>	<b>TONALITE (QUARTZ DIORITE) AND DIORITE</b>	Vermont Quartz Diorite— <i>biotite quartz diorite</i> (Griffith Park). Lar Quartz Diorite— <i>biotite quartz diorite, commonly foliated</i> (Griffith Park). Unnamed quartz diorites and diorite (Tehachapi Mtns., San Gabriel-Verdugo Mtns., Piru Creek area, San Emigdio Mtns., Santa Cruz Island); includes migmatized metasedimentary rocks in Tehachapi Mtns.
		bi	<b>MESOZOIC BASIC INTRUSIVE ROCKS</b>	Hornblende diorite and gabbro (Tehachapi Mtns., Verdugo Mtns.).
		ub	<b>MESOZOIC ULTRABASIC INTRUSIVE ROCKS</b>	Serpentine (San Rafael Mtns.).
		Ju	<b>UPPER JURASSIC MARINE SEDIMENTARY AND METASEDIMENTARY ROCKS</b>	Ju-Tr: Santa Monica Slate— <i>highly foliated dark-gray to black slate, with minor amounts of mica schist, phyllite, and spotted cordierite slate</i> (Late Jurassic pelecypods; formation probably Triassic in part, U.S. Geol. Survey Prof. Paper 420-A, p. 21-22).
		m	<b>PRE-CRETACEOUS METAMORPHIC ROCKS</b> <b>UNDIFFERENTIATED</b>	Pelona Schist— <i>highly foliated mica-chlorite-albite-quartz schists, some biotite schist, actinolite schist and quartzite; probably derived from sedimentary and pyroclastic rocks</i> (age uncertain, generally considered to be Precambrian or Mesozoic). Diorite gneiss and some biotite schist (San Gabriel Mtns.). Unnamed phyllite, chlorite schist and greenstone (Santa Cruz Island).
		ls	<b>ls = LIMESTONE AND/OR DOLOMITE</b>	Thin strata of light blue-gray fine crystalline limestone in the Pelona Schist. Lenses of marble in granitic and metamorphic rocks (Tehachapi Mtns. and area north of Mt. Pinos and Frazier Mtn.).
		ms	<b>PRE-CRETACEOUS METASEDIMENTARY ROCKS</b>	Placerita Formation— <i>graphite, sillimanite, and biotite schists, quartzite, and some limestone and dolomite</i> (San Gabriel Mtns.). Hornfels, quartzite, gneiss, marble, and schist (San Emigdio Mtns.). Biotite gneiss, minor mica schist, quartzite, and hornfels (Tehachapi Mtns.).
		mv	<b>PRE-CRETACEOUS METAVOLCANIC ROCKS</b>	Metamorphosed quartz latite (Soledad Mtn., Rosamond quadrangle).
JURASSIC		gr-m	<b>PRE-CENOZOIC GRANITIC AND METAMORPHIC ROCKS</b>	Diorite gneiss injected by granodiorite (migmatite) and in places includes the Placerita Formation (San Fernando quadrangle). Undifferentiated gneisses and granitic rocks (Mesozoic(?) to Precambrian; Mint Canyon area).
		IP	<b>PALEOZOIC MARINE SEDIMENTARY AND METASEDIMENTARY ROCKS</b>	Bean Canyon Formation— <i>dark-gray to black mica schist, phyllite, hornfels, and quartzite</i> (age uncertain, may be Mesozoic; formation in part volcanic—see IPv; Tehachapi Mtns.).
		ls	<b>ls = LIMESTONE AND/OR DOLOMITE</b>	Limestone, dolomite and marble of the Bean Canyon Formation.
		IPv	<b>PALEOZOIC METAVOLCANIC ROCKS</b>	Metabasalt and metafelsite flows or sills of the Bean Canyon Formation (age uncertain, may be Mesozoic; Tehachapi Mtns.).
		p-Can	<b>PRECAMBRIAN ANORTHOISITE</b>	Anorthosite <sup>10</sup> —Medium- to very coarse-grained; includes anorthosite intruded by Mesozoic(?) granite (San Gabriel Mtns.).
PALEOZOIC		p-Cgr	<b>UNDIVIDED PRECAMBRIAN GRANITIC ROCKS</b>	Echo Granite— <i>orange to pinkish-gray, locally foliated, quartz-rich granitic rocks</i> (San Gabriel Mtns.). Syenite phase of San Gabriel anorthosite complex (Mint Canyon area). Gabbro <sup>10</sup> — <i>gabbroic and noritic rocks; altered rocks bordering anorthosite; ilmenite-magnetite gabbro and massive ilmenite-magnetite</i> (San Gabriel Mtns.).
		p-Cg	<b>UNDIVIDED PRECAMBRIAN METAMORPHIC ROCKS:</b> <b>GNEISS</b>	Mendenhall Gneiss <sup>10</sup> — <i>quartz-plagioclase gneiss with ferromagnesian minerals</i> (San Gabriel Mtns.). Gneiss, veined gneiss, banded gneiss and migmatites (Frazier Mountain and Mt. Pinos areas). Coarse and fine augen gneisses, layered quartzofeldspathic gneisses, minor amphibolites (Mint Canyon area). Banded amphibolite, hornfels, diorite and granite gneiss (Sawtooth Mtns.). Foliated gneiss containing some recrystallized limestone and quartzite, local migmatite (Verdugo Mtns.). For some of these rocks the attributed Precambrian age is uncertain. Complex of gneiss and lesser amounts of granitic rocks, locally includes migmatites and dark hornblende diorites (the granitic rocks are of magmatic origin or are recrystallized from gneiss and often grade into gneiss; age Mesozoic or older), Sawmill Mountain-Bouquet Reservoir area.
		ls	<b>ls = LIMESTONE AND/OR DOLOMITE</b>	White coarsely crystalline marble lenses in gneiss, probably Precambrian, but possibly Paleozoic or Mesozoic (Sawmill Mountain area).

## NOTES

- <sup>1</sup> Late Pliocene and early Pleistocene age, W. O. Addicott, 1965, U.S. Geol. Survey Prof. Paper 503-B, p. B-8. Radiometric dates (K-Ar) from ash layer(s) near base of Santa Barbara Formation range from 11.5 ± 2.4 m.y. to 7.8 ± 1.2 m.y. reported by Robert S. Yeats and W. A. McLaughlin, Geol. Soc. America Special Paper, in press, 1968.
- <sup>2</sup> Probably Pleistocene, but may be late Pliocene in part.
- <sup>3</sup> May be Pleistocene (see G. T. James, 1963, Paleontology and nonmarine stratigraphy of the Cuyama Valley Badlands, California; Univ. Calif. Dept. Geol. Sci. Bull., vol. 45).
- <sup>4</sup> The Pico Formation includes rocks of early to late Pliocene age in the type locality in Ventura basin (Durham, D. L. and Yerkes, R. F., 1964, Geology and oil resources of the eastern Puente Hills area, southern California; U.S. Geol. Survey Prof. Paper 420-B, p. B-24-25); however, common usage in the L. A. basin restricts Pico to rocks of late Pliocene age. Modern U.S.G.S. usage assigns Pliocene marine rocks of the northeast L. A. basin to the upper member of the Fernando Formation.
- <sup>5</sup> "Repetto" is defined and properly used only as a stage designation. The U.S. Geological Survey has abandoned the name "Repetto Formation" and assigns these rocks to the lower member of the Fernando Formation (Durham and Yerkes—see ref. footnote 4).
- <sup>6</sup> Tequepis Sandstone and Sisquoc Formation are lateral equivalents; in part possibly of early Pliocene age in this map area.
- <sup>7</sup> Also called Maricopa Shale in San Emigdio Mountains and Modelo Formation in Ventura basin. Ranges in age from middle Miocene (Relizian) to late Miocene (Mohnian). In Cuyama Valley area divided into Whiterock Bluff Member, Saltos Shale Member and Branch Canyon Formation.
- <sup>8</sup> Oligocene or early Miocene age.
- <sup>9</sup> Radiometric age date (Pb<sup>206</sup>/U<sup>238</sup>) of 245 ± 10 m.y. determined by L. T. Silver, 1968, Preliminary history for the crystalline complex of the central Transverse Ranges, Los Angeles County, California; Geol. Soc. Amer. Spec. Paper no. 101 (Abstracts), p. 201-202. Also, oral communications from L. T. Silver, 1967 and 1968.
- <sup>10</sup> "Isotopic age determinations (U-Th-Pb) on zircons from pegmatite and granophyre dikes in the western San Gabriel Mountains have yielded concordant ages of about 1200 million years. From structural and petrological relations, this is both a minimum and a probable age for the San Gabriel Mountains anorthosite complex. It is also a minimum age for the Mendenhall Gneiss . . . which has been intruded by the anorthosite complex." L. T. Silver, et al., 1963, Precambrian age determinations in the western San Gabriel Mountains, California; Jour. Geology, vol. 71, no. 2, p. 196-214.

# INDEX TO GEOLOGIC MAPPING—Continued

40. Emery, Kenneth O., and Shepard, F. P., 1945, Lithology of the sea floor off southern California: *Geol. Soc. America Bull.*, v. 56, no. 4, pp. 431-478. (In addition, unpublished offshore geologic data collected by K. O. Emery, personal communication, 1961.)
- Vernon, James W. and Slater, R. A., 1963, Submarine tar mounds, Santa Barbara County, California: *Amer. Assoc. Petroleum Geologists Bull.*, v. 47, no. 8, pp. 1624-1627, fig. 1.
41. Frankian, R. T. and Associates, Report of Geologic and soil investigation, Verdugo Mountains development and conservation study, scale 1" = 400 feet, unpublished report for City of Glendale, 1968.
42. Fritsche, A. Eugene, Miocene geology of the central Sierra Madre Mountains, Santa Barbara County, California, scale 1:24,000, Univ. California, Los Angeles, Ph.D. thesis in progress, 1968.
43. Gamble, James H., Geology of the Point Mugu and Camarillo quadrangles, Ventura County, California, scale 1:12,000, Univ. California, Los Angeles, unpublished M.A. thesis, 1957.
44. Guynes, George E., Geology of the Triunfo Pass area, Los Angeles and Ventura Counties, scale 1:12,000, Univ. California, Los Angeles, unpublished M.A. thesis, 1959.
45. Hardey, Gordon W., Stratigraphy of the Topanga Formation in the eastern Santa Monica Mountains, scale 1:12,000, Univ. California, Los Angeles, unpublished M.A. thesis, 1958.
- Elliott, John L., Geology of the eastern Santa Monica Mountains, between Laurel Canyon and Beverly Glenn Boulevards, Los Angeles County, California, scale 1:12,000, Univ. California, Los Angeles, unpublished M.A. thesis, 1951.
46. Harding, Tod P., Geology of the eastern Santa Monica Mountains between Dry Canyon and Franklin Canyon, scale 1:10,000, Univ. California, Los Angeles, unpublished M.A. thesis, 1952.
47. Hetherington, George E., Geology of the south Topo Canyon area, Santa Susana quadrangle, Ventura County, California, scale 1:12,000, Univ. California, Los Angeles, unpublished M.A. thesis, 1957.
48. Hoots, Harold W., 1931, Geology of the eastern part of the Santa Monica Mountains, Los Angeles County, California: U. S. Geol. Survey Prof. Paper 165-C, 134 pp., Pl. 16, scale 1:24,000.
49. Jennings, Charles W., and T. W. Dibblee, Jr., Reconnaissance geologic mapping in parts of the Burnt Peak and Lake Hughes quadrangles, scale 1:24,000, California Div. Mines and Geology and U. S. Geol. Survey, 1967-68.
50. Jennings, Charles W. and Saul, Richard B., Reconnaissance geologic mapping and photo interpretation of part of the Burbank quadrangle, scale 1:24,000, California Div. Mines and Geology, reconnaissance mapping for the State Geologic Map, 1969.
51. Johnston, Robert L., The geology of a portion of the western Verdugo Mountains, scale approx. 1:12,000, Univ. California, Los Angeles, unpublished M.A. thesis, 1938.
52. Lamar, Donald L., Geology of the Elysian Park-Repetto Hills area, Los Angeles County, California, scale 1:24,000, report being prepared for publication by California Div. Mines and Geology, 1968.
- Lamar, Donald L., Structural evolution of the northern margin of the Los Angeles basin, scale 1:12,000, Univ. California, Los Angeles, unpublished Ph.D. thesis, 1961.
53. Levorsen, Robert I., Geology of the Las Lajas Canyon area, California, scale 1:7,200, Univ. California, Los Angeles, unpublished M.A. thesis, 1947.
54. Lofgren, Gary E., Geologic map of the Mt. Pinos basement complex, scale approx. 1:48,000, Stanford University, work in progress, 1968. Thrust faults from mapping by J. C. Crowell, 1964, ref. 18.
55. Martin, David R., Geology of the western part of the Santa Susana Mountains, Ventura County, California, scale 1:24,000, Univ. California, Los Angeles, unpublished M.A. thesis, 1958.
56. McGill, John T., Geology of a portion of the Las Flores and Dry Canyon quadrangles, Los Angeles County, California, scale 2 1/2 inches = approx. 1 mile, Univ. California, Los Angeles, unpublished M.A. thesis, 1948.
57. Metropolitan Water District of Southern California, unpublished geologic maps and sections, scale 1:12,000, prepared under the direction of R. J. Proctor:
- a) La Crescenta Tunnel route, L-1084 (including mapping by M.W.D. 1963-64, and from Robert L. Beatie, Geology of the Sunland-Tujunga area, Los Angeles County, California, scale approx. 1:12,000, Univ. California, Los Angeles, unpublished M.A. thesis, 1958).
- b) Verdugo Tunnel route, L-1079 (including mapping by M.W.D. 1964-66, and from Robert L. Beatie, see 57a).
- c) Flintridge Tunnel route, B-20263 (including mapping by M.W.D. 1964, and from J. P. Buwalda, unpublished report on Raymond basin to City of Pasadena, 1940).
- d) Altadena Tunnel route, B-20264, mapping by R. Crook, D. Kalin, M. Payne and R. Proctor of M.W.D. 1965-66.
- e) Santa Anita Tunnel No. 1 route, B-20265 (including mapping by M.W.D. 1964-66, and unpublished maps by P. Ehlig, California State College, Los Angeles, 1962, and D. Morton, California Div. Mines and Geology, 1966).
58. Metropolitan Water District of Southern California, unpublished geologic map and section, Sepulveda Tunnel area (L-1082), scale 1:12,000, prepared under the direction of R. J. Proctor, 1966-67. (In part after J. D. Traxler, 1948, see ref. 83.)
59. Morrison, Robert R., 1965, Map in Placerita-Soledad-Vasquez Rocks area: Pacific Section A.A.P.G. Guidebook, Map: The Geology of the Sand Canyon-Placerita Canyon area, scale approx. 1 inch = 3,100 feet.
60. Morton, Douglas M., Geologic map of the SE 1/4 of the Mt. Wilson quadrangle, scale 1:12,000, California Div. Mines and Geology, work in progress, 1968.
61. Muehlberger, William R., 1958, Geology of the northern Soledad basin, Los Angeles County, California: *Amer. Assoc. Petroleum Geol. Bull.* v. 42, no. 8, pp. 1812-1844, fig. 3, scale approx. 1:48,000.
- Jahns, Richard H. and Muehlberger, William R., 1954, Geology of the Soledad basin, Los Angeles County: *California Div. Mines Bull.* 170, Map sheet 6, scale approx. 1/4 inch = 1 mile.
62. Neuerburg, George J., 1953, Geology of the Griffith Park area, Los Angeles County, California: *California Div. Mines Special Rept.* 33, 29 pp., Pl. 1, scale approx. 3/8 inches = 1 mile. Modified in part by Woodford, et al (see ref. 91) and Durrell (see ref. 93).
63. Neuerburg, George J., Petrology of the pre-Cretaceous rocks of the Santa Monica Mountains, California, scale 1 inch = 400 feet, Univ. California, Los Angeles, unpublished Ph.D. thesis, 1951.
64. Oakshott, Gordon B., 1958, Geology and mineral deposits of San Fernando quadrangle, Los Angeles County, California: *California Div. Mines Bull.* 172, 147 pp., Pl. 1, scale 1:62,500.
65. Page, R. W., 1963, Geology and ground-water appraisal of the naval air missile test center area, Point Mugu, California: U. S. Geol. Survey Water-Supply Paper 1619-S, Pl. 1, scale 1:31,680 (addition of faults from mapping by Thomas Bailey, ref. 2).
66. Poland, Joseph F., Garrett, A. A., and Sinnott, Allen, 1959, Geology, hydrology, and chemical character of ground waters in the Torrance-Santa Monica area, California: U. S. Geol. Survey Water-Supply Paper 1461, Pl. 2, scale 1:31,680.
67. Pollard, Dalton L., Geology of the Halsey Canyon area, Los Angeles County, California, scale 1:12,000, Univ. California, Los Angeles, unpublished M.A. thesis, 1958.
68. Quarles, Miller, Jr., Geology of the Repetto and Montebello Hills, scale 1:24,000, Calif. Inst. Tech., unpublished M.S. thesis, 1941.
69. Renke, Daniel F., Geology of a part of Newbury Park quadrangle, Ventura County, California, scale 1:24,000, Univ. California, Los Angeles, unpublished M.A. thesis, 1957.
70. Robinson, Bobby B., Geology of the Holser Canyon area, Ventura County, California, scale 1:11,725, Univ. California, Los Angeles, unpublished M.A. thesis, 1956. (Subsurface extension of San Cayetano fault from Union Oil Co., letter 7/12/68.)
71. Rust, B. R., 1966, Late Cretaceous paleogeography near Wheeler Gorge, Ventura County, California: *Amer. Assoc. Petroleum Geol. Bull.* v. 50, no. 7, pp. 1389-1398, fig. 2, scale 1 inch = 3.5 miles.
72. Saul, Richard B., Geologic map of the SE 1/4 of the Oat Mountain quadrangle, scale 1:9,600, California Div. Mines and Geology work in progress, 1968.
73. Scholl, David W., 1960, Relationship of the insular shelf sediments to the sedimentary environments and geology of Anacapa Island, California: *Jour. Sedimentary Petrology*, v. 30, no. 1, pp. 123-139, fig. 3, scale 1:48,000.
74. Silver, Leon T., Basement geology of a part of the Soledad basin, scale 1:24,000, Calif. Inst. Tech. faculty research, 1966.
75. Sonneman, Howard S., Geology of the Boney Mountain area, Santa Monica Mountains, California, scale 1:12,000, Univ. California, Los Angeles, unpublished M.A. thesis, 1956.
76. Soper, E. K., 1938, Geology of the central Santa Monica Mountains, Los Angeles County: *California Div. Mines 34th Rept. State Mineralogist*, v. 34, no. 2, pp. 131-180, Pl. 1, scale 1:36,000.
77. Standard Oil Company of California, Western Operations, Inc., Geologic map of Santa Cruz Island, California, scale 1:48,000, unpublished.
78. Stanford Geological Survey under the direction of William R. Dickinson, Geology of the Pine Mountain area, Ventura County, California, scale 1:24,000, unpublished, 1964.
- Dickinson, William R. and Lowe, Donald R., 1966, Stratigraphic relations of phosphate- and gypsum-bearing upper Miocene strata, upper Sespe Creek, Ventura County, California: *Amer. Assoc. Petroleum Geol. Bull.* v. 50, no. 11, p. 2464-2481, fig. 2, scale 1 1/8" = 1 mile.
79. Stanley, Kenneth O., Structural history of the Clearwater fault, northwest Los Angeles County, California, scale 1:24,000, Univ. California, Los Angeles, unpublished M.A. thesis, 1966.
80. Superior Oil Company, Geologic map of Santa Cruz Island, by W. E. Kennett, scale 1 inch = 4000 feet, unpublished, 1944.
- Bremner, Carl St. J., 1932, Geology of Santa Cruz Island, Santa Barbara County, California: *Santa Barbara Mus. Nat. History Occ. Papers*, no. 1, 33 pp., Pl. 4, scale 1:62,500.
- Rand, William W., 1931, Preliminary report of the geology of Santa Cruz Island, Santa Barbara County, California: *California Div. Mines 27th Rept. State Mineralogist*, v. 27, no. 2, pp. 214-219: Geologic map of Santa Cruz Island, scale 1:62,500.
81. Szatai, John E., The geology of parts of Redrock Mountain, Warm Spring, Violin Canyon, and Red Mountain quadrangles, Los Angeles County, California, scale 1:24,000, Univ. Southern California, unpublished Ph.D. thesis, 1961. (Modification in the basal Castaic Formation from Calif. Dept. Water Resources, Geologic map of Castaic Reservoir, scale approx. 1:12,000.)
- Sams, Richard H., Geology of the Charlie Canyon area, northwest Los Angeles County, scale 1:9,600, Univ. California, Los Angeles, unpublished M.A. thesis, 1964.
- Konigsberg, Richard L., Geology along the San Francisquito fault, Los Angeles County, scale 1:12,000, Univ. California, Los Angeles, unpublished M.S. thesis, 1967.
82. Terpening, John N., Geology of a part of the eastern Santa Monica Mountains, scale 1:12,000, Univ. California, Los Angeles, unpublished M.A. thesis, 1951.
83. Traxler, J. Douglas, Geology of the east central Santa Monica Mountains, scale 1:24,000, Univ. California, Los Angeles, unpublished M.A. thesis, 1948.
84. Vedder, John G., 1968, Geologic map of Fox Mountain quadrangle, Santa Barbara County, California: U. S. Geol. Survey, Misc. Geol. Investigations Map I-547, scale 1:24,000.
85. Vedder, John G. and Brown, Robert D., Jr., Geologic map of parts of the San Rafael Mts., U. S. Geol. Survey, 1965 unpublished, and work in progress, 1968.
86. Vedder, John G. and Reppenning, Charles A., 1965, Geologic map of the southeastern Caliente Range, San Luis Obispo County, California: U. S. Geol. Survey Oil and Gas Investigations Map OM-217, scale 1:24,000.
87. Vedder, J. G., Gower, H. D., Clifton, H. E., and Durham, D. L., 1967, Reconnaissance geologic map of the central San Rafael Mountains and vicinity, Santa Barbara County, California: U. S. Geol. Survey Miscellaneous Geologic Investigations Map I-487, scale 1:48,000 (map by authors and T. W. Dibblee, Jr. and E. E. Brabb).
88. Weber, F. Harold, Jr., Geologic map of part of the Thousand Oaks quadrangle, scale 1:9,600, California Div. Mines and Geology, work in progress, 1968.
89. Wiese, John H., 1950, Geology and mineral resources of the Neenach quadrangle, California: *California Div. Mines Bull.* 153, 53 pp., Pl. 1, scale 1:62,500. (Differentiation of marble from other metasedimentary rocks and certain fault additions by T. W. Dibblee, Jr. unpublished geologic maps, 1957, scale 1:24,000.)
90. Winterer, E. L. and Durham, D. L., 1962, Geology of southeastern Ventura basin, Los Angeles County, California: U. S. Geol. Survey Prof. Paper 334-H, 366 pp., Pl. 44, scale 1:24,000.
91. Woodford, A. O., Schoellhamer, J. E., Vedder, J. G., and Yerkes, R. F., 1954, Geology of the Los Angeles basin: *in California Div. Mines Bull.* 170, chapt. II, contr. 5, pp. 65-83, Pl. 1, scale 1:96,000.
92. Yeats, Robert S., 1965, Pliocene seaknoll at South Mountain, Ventura basin, California: *Amer. Association Petroleum Geol. Bull.* v. 49, no. 5, pp. 526-546, fig. 2, scale 1 1/8" = 1 mile.
93. Yerkes, R. F., Campbell, R. H., Schoellhamer, J. E., Wentworth, C. M. and Blackerby, B. A., Geologic maps of the central Santa Monica Mountains, U. S. Geol. Survey open file and manuscript maps, scale 1:12,000, 1968. (For another structural interpretation of this area, see Cordell Durrell, 1954, Geology of the Santa Monica Mountains, Los Angeles and Ventura Counties: *California Div. Mines Bull.* 170, Map sheet 8, scale approx. 1:125,000.)

For a complete list of published geologic maps of this area see Division of Mines and Geology Special Reports 52, 52-A and 52-B.



The San Andreas, Garlock, and Big Pine faults and their intersections are the prominent features of this photo that portrays most of the eastern half of the Los Angeles Sheet area. View is southeasterly from approximately 65,000 feet above Interstate 5 near the northern edge of the map sheet area. Photo of November, 1967, courtesy U. S. Geological Survey.